



PHENIX measurements of single electrons from charm and bottom decays at midrapidity in Au+Au collisions

Kazuya Nagashima


for the PHENIX Collaboration
(Hiroshima University/RIKEN)

✓ Heavy Flavor Suppression

PHENIX measured strong suppression of $\text{HF} \rightarrow e$

Heavy flavor energy loss

- Collisional energy loss \sim low p_T
 - > Langevin approach
- Radiative energy loss \sim high p_T
 - > Bethe-Heitler formula
 - > Dead cone effect



Mass ordering (expected)

$$\Delta E_g > \Delta E_{u,d,s} > \Delta E_c > \Delta E_b \quad \rightarrow \quad R_{AA}(u,d,s) < R_{AA}(c) < R_{AA}(b) ?$$

✓ Charm and Bottom Suppression

PHENIX found

$$R_{AA}(b \rightarrow e) > R_{AA}(c \rightarrow e)$$

[In 2014-2016]

PHENIX collected


AuAu (HF): 4.9nb^{-1}

pp: 23pb^{-1}



- > high precision
- > centrality dependence
- > flow measurement

provide strong constraint
on QGP property

Today's result use 1/8 of full dataset





✓ PHENIX Silicon Vertex Detector (VTX)



Silicon Vertex Detector (VTX)

- installed in 2011
- 2 pixel layers + 2 strip layers
 $(\sigma_\phi = 14.4 \mu\text{m})$ $(\sigma_\phi = 23 \mu\text{m})$
- precise collision vertex

✓ PHENIX Silicon Vertex Detector (VTX)



Silicon Vertex Detector (VTX)

- installed in 2011
- 2 pixel layers + 2 strip layers
 $(\sigma_\phi = 14.4 \mu\text{m})$ $(\sigma_\phi = 23 \mu\text{m})$
- precise collision vertex

Precise displaced tracking


- Distance of Closest Approach(DCA)
 $\mathbf{DCA}_T = \mathbf{L} - \mathbf{R}$
- DCA analysis allows a separation of
 $B^0 \rightarrow e$ from $D^0 \rightarrow e$
 $B^0 : c\tau = 455.4 \mu\text{m}$
 $D^0 : c\tau = 122.9 \mu\text{m}$
- almost all BG from $\mathbf{DCA}_T = 0$

✓ Bayesian Inference Techniques

- Bayes' theorem $P(\theta|x) \propto P(x|\theta)\pi(\theta)$
- Simultaneous fit to dN/dp_T and $DCA_T(p_T)$



Measured data
 $dN/dp_T, DCA_T(p_T)$

$P(x|\theta)\pi(\theta)$
Likelihood




✓ Bayesian Inference Techniques

- Bayes' theorem $P(\theta|x) \propto P(x|\theta)\pi(\theta)$
- Simultaneous fit to dN/dp_T and $DCA_T(p_T)$





✓ Bayesian Inference Techniques

- Bayes' theorem $P(\theta|x) \propto P(x|\theta)\pi(\theta)$
- Simultaneous fit to dN/dp_T and $DCA_T(p_T)$





✓ Bayesian Inference Techniques

- Bayes' theorem $P(\theta|x) \propto P(x|\theta)\pi(\theta)$
- Simultaneous fit to dN/dp_T and $DCA_T(p_T)$
- employ Markov Chain Monte Carlo (MCMC) for sampling





✓ Bayesian Inference Techniques

- Bayes' theorem $P(\theta|x) \propto P(x|\theta)\pi(\theta)$
- Simultaneous fit to dN/dp_T and $DCA_T(p_T)$
- employ Markov Chain Monte Carlo (MCMC) for sampling







Comparison to Data in 0-10% central



Simultaneous fit to Invariant yield and DCA_T(p_T)




Comparison to Data in 0-10% central



Simultaneous fit to Invariant yield and DCA_T(p_T)
Unfolding results agree well with measured data



✓ Comparison to Data in 0-10% central



- Invariant yields of **charm** and **bottom** hadrons



Comparison to Data in 0-10% central




- Invariant yields of **charm** and **bottom** hadrons
- D⁰ extracted from c hadrons and PYTHIA model
→ agree with the STAR direct measurement

✓ Bottom Electron Fraction

PHENIX published bottom electron fraction in MB AuAu
(*Phys. Rev. C, 93, 034904*)

New result reproduce published result well




✓ Bottom Electron Fraction

PHENIX published bottom electron fraction in MB AuAu
(*Phys. Rev. C, 93, 034904*)

New

New result reproduce published result well

b-fraction in 0-10% AuAu seems higher than MB
(consistent within uncertainty)
→ need more precision



✓ Charm and Bottom Suppression

R_{AA} of c → e and b → e

$$R_{AA}^{b \rightarrow e} = \frac{F_{AuAu}}{F_{pp}} R_{AA}^{HF}$$

$$R_{AA}^{c \rightarrow e} = \frac{(1 - F_{AuAu})}{(1 - F_{pp})} R_{AA}^{HF}$$


R_{AA}^{HF} : PHENIX measurement

F_{AuAu} : Unfolding result
(1/4 of 2014 data)

F_{pp} : STAR e-h correlations
(p_T : 3.0~8.0 GeV/c)

2004+2011 AuAu MB

Phys.Rev.C, 93, 034904



✓ Charm and Bottom Suppression

R_{AA} of $c \rightarrow e$ and $b \rightarrow e$

$$R_{AA}^{b \rightarrow e} = \frac{F_{AuAu}}{F_{pp}} R_{AA}^{HF}$$

$$R_{AA}^{c \rightarrow e} = \frac{(1 - F_{AuAu})}{(1 - F_{pp})} R_{AA}^{HF}$$


R_{AA}^{HF} : PHENIX measurement

F_{AuAu} : Unfolding result
(1/4 of 2014 data)



F_{pp} : STAR e-h correlations
(p_T : 3.0~8.0 GeV/c)

New result reproduce well
with published result

2004+2014 AuAu MB





✓ R_{AA} of $c \rightarrow e$ and $b \rightarrow e$



> $b \rightarrow e$ is less suppressed than $c \rightarrow e$ in 3.0-5.0 GeV/c (0-10% AuAu)

✓ R_{AA} of $c \rightarrow e$ and $b \rightarrow e$




- > $b \rightarrow e$ is less suppressed than $c \rightarrow e$ in 3.0-5.0 GeV/c (0-10% AuAu)
- > $c \rightarrow e$ in 0-10% is suppressed stronger than $c \rightarrow e$ in MB
- Other centrality results are ongoing

✓ Model Comparison in Bottom Fraction

[Theoretical models]

T-Matrix approach

- > hadron resonance in QGP
- > Diffusion coefficient: 4-30
- $D(2\pi T) < 4$
- strong coupling in QGP



✓ Model Comparison in Bottom Fraction


[Theoretical models]

T-Matrix approach

- > hadron resonance in QGP
- > Diffusion coefficient: 4-30
- $D(2\pi T) < 4$
- strong coupling in QGP

DGLV model

- > rad. + coll. energy loss
- > dynamical E-loss formalism
- reasonable agreement




✓ Model Comparison in R_{AA}

0-10% central

[Theoretical models]

DGLV model

> coll. + rad. (dynamical)



✓ Model Comparison in R_{AA}

0-10% central

[Theoretical models]


DGLV model

> coll. + rad. (dynamical)

SUBATECH

> coll. + rad.(LPM)

> Hard Thermal Loop



✓ Model Comparison in R_{AA}

0-10% central

[Theoretical models]

DGLV model

> coll. + rad. (dynamical)


SUBATECH

> coll. + rad.(LPM)

> Hard Thermal Loop

T-Matrix approach

> hadron resonance in QGP
(parameter free approach)



✓ Model Comparison in R_{AA}

0-10% central

[Theoretical models]

DGLV model

> coll. + rad. (dynamical)


SUBATECH

> coll. + rad.(LPM)

> Hard Thermal Loop

T-Matrix approach

> hadron resonance in QGP
(parameter free approach)



Theory: reasonable agreement, but need smaller diffusion?


Data: smaller uncertainty and broader p_T range (coming soon)

✓ Future Prospects

New

[Update points]

- use full statistics ($\times 8$)
- new inclusive dN/dp_T
- new pp reference
 - > smaller systematic uncertainty
 - > broader p_T range (1-9 GeV/c) for R_{AA} of $c \rightarrow e$, $b \rightarrow e$
 - > R_{AA} of c and b hadrons
- PHENIX also measure $B \rightarrow J/\psi$
 - > very low p_T B meson
 - see Cesar Silva's Talk (8.4 – Wed 16:30)



✓ Summary

- Measurement of $R_{AA}^{c \rightarrow e}$ and $R_{AA}^{b \rightarrow e}$ in 0-10% AuAu

- $R_{AA}^{b \rightarrow e} > R_{AA}^{c \rightarrow e}$ (in 3.0-5.0 GeV/c)
- $R_{AA}^{c \rightarrow e}$ (MB) > $R_{AA}^{c \rightarrow e}$ (0-10%)

- Model Comparison

Bottom electron fraction

> strong coupling in QGP ($D(2\pi T) < 4$)

Nuclear modification factor R_{AA}

> reasonable agreement, but needs smaller diffusion?

- Future prospects

- Full dataset in AuAu and pp provide more precise
- Measurement of $R_{AA}^{b \rightarrow e}$ and $R_{AA}^{c \rightarrow e}$
in broader p_T range ~ 1.0 - 9.0 GeV/c

→ Final result is coming soon!!

Thank you !!

also see

B \rightarrow J/ ψ at forward
(Cesar Luiz da Silva, 8.4 – Wed 16:30)
HF in small system
(Sanghoon Lim, 8.4 – Wed 17:10)




Backup




Bottom Electron Fraction

Phys. Rev. C, 93, 034904



✓ Testing Possible Baryon Enhancement



- Follow P. Sorensen and X. Dong
(*Phys Rev C 74, 024902 (2006)*)
- Λ/K_s ratio measured
in STAR 20-40% Au+Au at 200 GeV
and STAR in p+p at 200 GeV
(*arXiv:nucl-ex/0601042*)
- Fit both data
- Fix asymptotic value to 0.3
in both Au +Au and p+p
- Apply enhanced the ratio of
 Λ_c/D and Λ_b/B to the decay matrices.






Baryon Enhancement

Talk by Long Zhou (@QM2017)



Charm baryon enhancement is similar
to strangeness baryon enhancement

✓ Testing Possible Baryon Enhancement





b-fraction increase when assuming enhanced baryon.

Covered by systematic uncertainty of the measurement

Not included additional uncertainty



Testing Possible Baryon Enhancement




Covered by systematic uncertainty of the measurement
Not included additional uncertainty



New DCA_T Distribution in 0-10% AuAu

$2.00 < p_T < 2.50$




DCA_T distribution of electrons


- $1.25 < p_T < 6.0$ GeV (9 binning)
- not corrected for PHENIX acceptance and efficiency

DCA_T resolution

- $\sigma = 0.006$ cm @ $p_T = 2.5$ GeV/c



✓ New DCA_T Distribution in 0-10% AuAu









BG components

- Mis-reconstruction
 - > Mis-ID hadron
 - > High-multiplicity
- Prompt
 - > Dalitz, J/ ψ
- Non-prompt
 - > Conversion, K_{e3}

Heavy flavor decay electron

- dominates at $|0.02| < \text{DCA}_T < |0.1|$

✓ MC Sampling for



- assuming Gaussian uncertainties of F_{AuAu} , F_{pp} , R_{AA}
- median and error get from MC sampling

$$R_{AA}^{b \rightarrow e} = \frac{F_{AuAu}}{F_{pp}} R_{AA}^{HF}$$

$$R_{AA}^{c \rightarrow e} = \frac{(1 - F_{AuAu})}{(1 - F_{pp})} R_{AA}^{HF}$$

✓ Background Normalization

Non-photonic to photonic electron ratio R_{NP}

$$R_{NP} = N_{NP} / N_P$$

$$N_P = (N_e^{\text{iso}} - \varepsilon_R * \varepsilon_I * N_e) / \varepsilon_R * (1 - \varepsilon_I)$$

$$N_{NP} = (\varepsilon_R * N_e^{\text{iso}} - N_e) / \varepsilon_R * (1 - \varepsilon_I)$$

$$N_e = N_{NP} + N_P$$

$$N_e^{\text{iso}} = \varepsilon_R * N_{NP} + \varepsilon_R * \varepsilon_I * N_P$$


(applied isolation cut)

ε_I = single cut efficiency

(calculated by MC method)

ε_R = randomly killed efficiency

(calculated by data driven)



✓ Background Normalization



Track swapping method

[Mis-ID hadron]

- Swapped at RICH
- opposite side RICH hit


[High-multiplicity]

- Swapped at VTX
- 10 degree rotation







Systematic Uncertainty of b-fraction







Decay Probability





Comparison to Data in minimum bias




Unfolding results agree with measured data well




Comparison to Data

Minimum bias




0-10% central




Invariant yields of **charm** and **bottom** hadrons

✓ Comparison to Measured Data



✓ Full probability distribution of parameters



[Diagonal]
Sampled yield

[Other]
Correlation between
c and b hadron yield

✓ Energy loss model

collisional energy loss

- parton elastic scattering
- Brownian motion via Langevin equation

$$\frac{d\vec{p}}{dt} = -\eta_D(p)\vec{p} + \vec{\xi}$$
 η_D : friction coefficient
 $\vec{\xi}$: drift force

radiative energy loss

- Bathe-Heitler for gluon radiation

$$dP_0 \approx \frac{\alpha_s C_F}{\pi} \frac{d\omega}{\omega} \frac{dk_\perp^2}{k_\perp^2}$$

> Dead-Cone effect

- strong suppression of HF in small-angle radiation

$$\propto \frac{k_\perp^2 dk_\perp^2}{(k_\perp^2 + \omega^2 \theta_0^2)^2}, (\theta_0 \equiv \frac{M}{E})$$


> Landau-Pomeranchuk-Migdal effect

- suppression in high density $\propto \frac{\lambda_{path}}{L_{form}}$


mass ordering

$$\Delta E_g > \Delta E_{u,d,s} > (?) \Delta E_c > (?) \Delta E_b$$

Dead-Cone effect



LPM effect



medium

λ_{path} : mean free path

L_{form} : formation length

